Documentation and Reporting Heart Failure

DOCUMENTATION TIPS

Document and report heart failure to the highest level of specificity known for each visit. Heart failure cannot be reported from diagnostic tests alone. The provider must bring the test results into the progress note and report the diagnosis including status and treatment plan. Documentation of diastolic or systolic **dysfunction** does not support reporting heart failure. Documentation should identify:

- Location left ventricular, right-sided, biventricular, systolic, diastolic or combined
- Chronicity acute, chronic, acute on chronic
- **Underlying conditions and/or causal relationships** hypertension, diabetes, coronary artery disease, arrhythmia, kidney disease, obesity, congenital heart disease

Document clinical indicators in the absence of heart failure:

- Non-exertional dyspnea
- Edema

Fatigue

• Weight gain

Heart failure (not an all-inclusive list)

ICD-10-CM Code	Description
150.1	Left ventricular failure, unspecified
150.2-	Systolic (congestive) heart failure
150.3-	Diastolic (congestive) heart failure
150.4-	Combined systolic (congestive) and diastolic (congestive) heart failure
I50.81-	Right heart failure
150.82	Biventricular heart failure
150.83	High output heart failure
150.84	End stage heart failure
150.9	Heart failure (congestive), unspecified

Hypertensive heart disease and chronic kidney disease (not an all-inclusive list)

ICD-10-CM Code	Description
111.0	Hypertensive heart disease with heart failure
111.9	Hypertensive heart disease without heart failure
I13.0	Hypertensive heart and chronic kidney disease with heart failure and stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease



ICD-10-CM Code	Description
l13.10	Hypertensive heart and chronic kidney disease without heart failure and stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease
l13.11	Hypertensive heart and chronic kidney disease without heart failure, with stage 5 chronic kidney disease, or end stage renal disease
113.2	Hypertensive heart and chronic kidney disease with heart failure, with stage 5 chronic kidney disease, or end stage renal disease

Report the type of heart failure (I50.-) and the stage of chronic kidney disease (N18.-), as applicable.

Document and report any tobacco exposure, dependence or tobacco dependence in remission.

DOCUMENTATION AND REPORTING EXAMPLE

Patient with hypertension, CKD, stage 3b, and chronic combined systolic and diastolic heart failure. Continue Toprol 100 mg b.i.d. for heart failure. BP 145/80 – continue Lisinopril, 10 mg q.d. Labs from 7/10/2021, eGFR is 34, follow-up with nephrologist.

I13.0 Hypertensive heart and chronic kidney disease with heart failure and stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease N18.32 Chronic kidney disease stage 3b

150.42 Chronic combined systolic and diastolic heart failure



HEDIS: Detailed information about measures related to hypertension or kidney disease is available through your Passport/Molina Quality Representative.

According to the ICD-10-CM Official Guidelines for Coding and Reporting FY2022: "A dash (-) at the end of an alphabetic index entry indicates that additional characters are required." Refer to the tabular list to identify the appropriate character(s) that will complete the diagnosis code.



ICD-10-CM Official Guidelines for Coding and Reporting FY2022. 1 Oct. 2021, www.cms.gov/files/document/fy-2022-icd-10-cm-coding-guidelines.pdf. Accessed 30 Nov. 2021.

^{2.} Watson, R D et al. "ABC of heart failure. Clinical features and complications." BMJ (Clinical research ed.) vol. 320,7229 (2000): 236-9. doi:10.1136/bmj.320.7229.236

Malik A, Brito D, Chhabra L. "Heart Failure: Types, Symptoms, Causes & Treatments." Cleveland Clinic, StatPearls Publishing, January 2020. https://my.clevelandclinic.org/health/diseases/17069-heart-failure-understanding-heart-failure